

On the existence and uniqueness of minima and maxima on spheres of the integral functional of the calculus of variations

BIAGIO RICCERI

Abstract: Given a bounded domain $\Omega \subset \mathbf{R}^n$, we prove that if $f : \mathbf{R}^{n+1} \rightarrow \mathbf{R}$ is a C^1 function whose gradient is Lipschitzian in $\mathbf{R}^{n+1}$ and non-zero at 0, then, for each $r > 0$ small enough, the restriction of the integral functional $u \rightarrow \int_{\Omega} f(u(x), \nabla u(x)) dx$ to the sphere $\{u \in H^1(\Omega) : \int_{\Omega} (|\nabla u(x)|^2 + |u(x)|^2) dx = r\}$ has a unique global minimum and a unique global maximum.

Key words: Sobolev space; integral functional; minimum; maximum; sphere; existence; uniqueness.

2000 Mathematics Subject Classification. 49K27.

Introduction

Here and in the sequel, $\Omega \subset \mathbf{R}^n$ is a bounded domain, and $f : \mathbf{R}^{n+1} \rightarrow \mathbf{R}$ is a C^1 function whose gradient is non-constant and Lipschitzian (with respect to the Euclidean metric).

We will consider the Sobolev space $H^1(\Omega)$ endowed with the norm

$$\|u\| = \left(\int_{\Omega} (|\nabla u(x)|^2 + |u(x)|^2) dx \right)^{\frac{1}{2}}$$

which is induced by the scalar product

$$\langle u, v \rangle = \int_{\Omega} (\nabla u(x) \nabla v(x) + u(x) v(x)) dx .$$

The linear growth of ∇f (coming from its Lipschitzianity) implies that the functional

$$u \rightarrow J(u) := \int_{\Omega} f(u(x), \nabla u(x)) dx$$

is (well-defined and) C^1 on $H^1(\Omega)$, with derivative given by

$$\langle J'(u), v \rangle = \int_{\Omega} (f_{\xi}(u(x), \nabla u(x)) v(x) + \nabla_{\eta} f(u(x), \nabla u(x)) \nabla v(x)) dx$$

for all $u, v \in H^1(\Omega)$ ([2], p. 249).

Let $r > 0$. We are interested in minima and maxima of the restriction of the functional J to the sphere $S_r := \{u \in H^1(\Omega) : \|u\| = r\}$.

In the present setting, there is no evidence of their existence and uniqueness. In fact, with regard to the existence aspect, not only S_r is not weakly compact but also, if $f(\xi, \cdot)$ is neither convex nor concave in $\mathbf{R}^n$, the functional J is neither lower nor upper weakly semicontinuous. But, even when J is sequentially weakly continuous, it may happen that J has no minima and/or maxima on S_r .

In this connection, consider the following simple and enlightening situation. Assume that f depends only on the first variable and that has a unique global maximum in $\mathbf{R}$, say ξ_0 . So, $J(u) = \int_{\Omega} f(u(x))dx$. Then, it is clear that the constant function $x \rightarrow \xi_0$ is the unique maximum of the functional J . In this case, J turns out to be sequentially weakly continuous, thanks to the Rellich-Kondrachov theorem. Then, by Lemma 2.1 of [1], the function $\rho \rightarrow \sup_{S_{\rho}} J$ is non-decreasing in $]0, +\infty[$. Consequently, if $r > |\xi_0|(\text{meas}(\Omega))^{\frac{1}{2}}$, $J|_{S_r}$ has no maxima.

Nevertheless, we will show that if $\nabla f(0) \neq 0$ then $J|_{S_r}$ possesses exactly one minimum and exactly one maximum for each $r > 0$ small enough.

The result

To shorten the statement of our result, let us introduce some further notations. In the sequel, $g : \mathbf{R}^{n+1} \rightarrow \mathbf{R}$ is another C^1 function which is non-negative, with $g(0) = 0$, and whose gradient is Lipschitzian, with Lipschitz constant $\nu < 2$. We set

$$I(u) = \int_{\Omega} g(u(x), \nabla u(x))dx$$

for all $u \in H^1(\Omega)$.

Moreover, V is a closed linear subspace of $H^1(\Omega)$ with the following property: there exists $v_0 \in V$ such that

$$\int_{\Omega} (f_{\xi}(0)v_0(x) + \nabla_{\eta} f(0)\nabla v_0(x))dx \neq 0 .$$

Finally, if L is the Lipschitz constant of ∇f , we denote by S the set (possibly empty) of all global minima of the restriction to V of the functional

$$u \rightarrow \|u\|^2 + I(u) + \frac{2-\nu}{L}J(u) .$$

Then, with the convention $\inf \emptyset = +\infty$, our result reads as follows:

THEOREM 1. - Under the above assumptions, one has

$$\delta := \inf_{u \in S} (\|u\|^2 + I(u)) > 0$$

and, for each $r \in]0, \delta[$, the restriction of the functional J to the set

$$C_r := \{u \in V : \|u\|^2 + I(u) = r\}$$

has a unique global minimum.

PROOF. Let $\mu \geq 0$ and let $u, v, w \in H^1(\Omega)$, with $\|w\| = 1$. Using Cauchy-Schwartz and Hölder inequalities, we have

$$\begin{aligned} & |\langle I'(u) + \mu J'(u) - I'(v) - \mu J'(v), w \rangle| \leq \\ & \leq \int_{\Omega} |(g_{\xi}(u, \nabla u) - g_{\xi}(v, \nabla v))w + (\nabla_{\eta} g(u, \nabla u) - \nabla_{\eta} g(v, \nabla v)) \nabla w| dx + \\ & + \mu \int_{\Omega} |(f_{\xi}(u, \nabla u) - f_{\xi}(v, \nabla v))w + (\nabla_{\eta} f(u, \nabla u) - \nabla_{\eta} f(v, \nabla v)) \nabla w| dx \leq \\ & \leq \int_{\Omega} (|g_{\xi}(u, \nabla u) - g_{\xi}(v, \nabla v)|^2 + |\nabla_{\eta} g(u, \nabla u) - \nabla_{\eta} g(v, \nabla v)|^2)^{\frac{1}{2}} (|w|^2 + |\nabla w|^2)^{\frac{1}{2}} dx + \\ & + \mu \int_{\Omega} (|f_{\xi}(u, \nabla u) - f_{\xi}(v, \nabla v)|^2 + |\nabla_{\eta} f(u, \nabla u) - \nabla_{\eta} f(v, \nabla v)|^2)^{\frac{1}{2}} (|w|^2 + |\nabla w|^2)^{\frac{1}{2}} dx \leq \\ & \leq \left(\int_{\Omega} (|g_{\xi}(u, \nabla u) - g_{\xi}(v, \nabla v)|^2 + |\nabla_{\eta} g(u, \nabla u) - \nabla_{\eta} g(v, \nabla v)|^2) dx \right)^{\frac{1}{2}} + \\ & + \mu \left(\int_{\Omega} (|f_{\xi}(u, \nabla u) - f_{\xi}(v, \nabla v)|^2 + |\nabla_{\eta} f(u, \nabla u) - \nabla_{\eta} f(v, \nabla v)|^2) dx \right)^{\frac{1}{2}} \leq \\ & \leq (\nu + \mu L) \|u - v\|. \end{aligned}$$

Hence, the derivative of the functional $I + \mu J$ is Lipschitzian, with constant $\nu + \mu L$. As a consequence, if $0 \leq \mu < \frac{2-\nu}{L}$, the functional $u \rightarrow \|u\|^2 + I(u) + \mu J(u)$ is strictly convex and coercive. To see this, it is enough to show that its derivative is strongly monotone ([3], pp. 247-248). Indeed, if $\Phi(\cdot) := \|\cdot\|^2$, we have for all $u, v \in H^1(\Omega)$

$$\begin{aligned} & \langle \Phi'(u) + I'(u) + \mu J'(u) - \Phi'(v) - I'(v) - \mu J'(v), u - v \rangle \geq \\ & \geq 2\|u - v\|^2 - \|I'(u) - J'(v) + \mu(I'(v) - J'(v))\| \|u - v\| \geq (2 - \nu - \mu L) \|u - v\|^2. \end{aligned}$$

Clearly, this shows also the convexity of the functional $\Phi + I + \frac{2-\nu}{L} J$. Assume $S \neq \emptyset$. Then, S is closed and convex, and so there exists a unique $\hat{u} \in S$ such that

$$\|\hat{u}\|^2 + I(\hat{u}) = \delta.$$

Observe that $\|u\|^2 + I(u) > 0$ for all $u \in V \setminus \{0\}$. So, $\delta \geq 0$. Arguing by contradiction, assume $\delta = 0$. Then, it would follow $\hat{u} = 0$. Hence, since $0 \in S$, we would have

$$\langle \Phi'(0) + I'(0) + \frac{2-\nu}{L}J'(0), v \rangle = 0$$

for all $u \in V$ and so, since $\Phi'(0) + I'(0) = 0$ (being 0 the global minimum of $\Phi + I$), it would follow

$$\int_{\Omega} (f_{\xi}(0)v(x) + \nabla_{\eta}f(0)\nabla v(x))dx = 0$$

for all $v \in V$, against one of the hypotheses. Hence, we have proven that $\delta > 0$. Now, fix $r \in]0, \delta[$ and consider the function $\Psi : V \times [\frac{L}{2-\nu}, +\infty[\rightarrow \mathbf{R}$ defined by

$$\Psi(u, \lambda) = J(u) + \lambda(\|u\|^2 + I(u) - r)$$

for all $(u, \lambda) \in V \times [\frac{L}{2-\nu}, +\infty[$. As we have seen above, $\Psi(\cdot, \lambda)$ is continuous and convex for all $\lambda \geq \frac{L}{2-\nu}$ and coercive for all $\lambda > \frac{L}{2-\nu}$, while $\Psi(u, \cdot)$ is continuous and concave for all $u \in V$, with $\lim_{\lambda \rightarrow +\infty} \Psi(0, \lambda) = -\infty$. So, we can apply to Ψ a classical saddle-point theorem ([3], Theorem 49.A) which ensures the existence of $(u^*, \lambda^*) \in V \times [\frac{L}{2-\nu}, +\infty[$ such that

$$\begin{aligned} J(u^*) + \lambda^*(\|u^*\|^2 + I(u^*) - r) &= \inf_{u \in V} (J(u) + \lambda^*(\|u\|^2 + I(u) - r)) = \\ &= J(u^*) + \sup_{\lambda \geq \frac{L}{2-\nu}} \lambda(\|u^*\|^2 + I(u^*) - r) . \end{aligned}$$

Of course, we have $\|u^*\|^2 + I(u^*) \leq r$, since the sup is finite. But, if it were $\|u^*\|^2 + I(u^*) < r$, we would have $\lambda^* = \frac{L}{2-\nu}$. This, in turn, would imply that $u^* \in S$, against the fact that $r < \delta$. Hence, we have $\|u^*\|^2 + I(u^*) = r$. Consequently

$$J(u^*) + \lambda^*r = \inf_{u \in V} (J(u) + \lambda^*(\|u\|^2 + I(u))) .$$

From this, we infer that $\lambda^* > \frac{L}{2-\nu}$ (since $r < \delta$), that u^* is a global minimum of $J|_{C_r}$ and that if each global minimum of $J|_{C_r}$ is a global minimum in V of the functional $u \rightarrow \|u\|^2 + I(u) + \lambda^*J(u)$. Since $\lambda^* > \frac{L}{2-\nu}$, this functional is strictly convex and so u^* is its unique global minimum in V . The proof is complete. $\triangle$

REMARK 1. It is almost superfluous to remark that the conclusion of Theorem 1 may fail if the assumption that involves V and $\nabla f(0)$ is not satisfied. In this connection, consider, for instance, the case $f(\sigma) = -|\sigma|^2$, with $g = 0$. This assumption, however, serves only to ensure that $\delta > 0$. So, it becomes superfluous, in particular, when $S = \emptyset$.

Now, denote by S_1 the set (possibly empty) of all global minima of the restriction to V of the functional

$$u \rightarrow \|u\|^2 + I(u) - \frac{2-\nu}{L}J(u) .$$

Clearly, applying Theorem 1 also to $-f$, we get

THEOREM 2. - *Under the assumptions of Theorem 1, one has*

$$\delta_1 := \min \left\{ \inf_{u \in S} (\|u\|^2 + I(u)), \inf_{u \in S_1} (\|u\|^2 + I(u)) \right\} > 0$$

and, for each $r \in]0, \delta_1[$, the restriction of the functional J to the set

$$\{u \in V : \|u\|^2 + I(u) = r\}$$

has a unique global minimum and a unique global maximum.

References

- [1] M. SCHECHTER and K. TINTAREV, *Spherical maxima in Hilbert space and semi-linear elliptic eigenvalue problems*, Differential Integral Equations, **3** (1990), 889-899.
- [2] M. STRUWE, *Variational methods*, Springer-Verlag, 1996.
- [3] E. ZEIDLER, *Nonlinear functional analysis and its applications*, vol. III, Springer-Verlag, 1985.

Department of Mathematics
University of Catania
Viale A. Doria 6
95125 Catania
Italy
e-mail address: ricceri@dmf.unict.it